

## **Crimson Clover** (*Trifolium incarnatum* L.)



### Varieties:

**AU Robin, Chief, Dixie, Fame, Tibbee**

\*Grateful Acknowledgment for permission to use the photo which is copyrighted by International Plant Nutrition Institute©

### Growth Habit:

**Cool Season Annual**

### Area of Adaptation:

**North, Central, South**

### Soil Type:

**Well drained**

### pH Range:

**6.0-7.0**

### Seeding Rate (lb/A) PLS:

**B: 20-30**

### Seeding Depth (in):

**1/4-1/2**

### Seed Quality (%Germ, Purity, Weed)

**85 99 0.50**

### Seed/Pound:

**150,000**

### Seedling Vigor:

**Good**

### Inoculant Group:

**Clover**

### Planting Date:

**N: Aug. 25 – Oct. 1**

**C: Sept. 1 – Oct. 15**

**S: Sept. 15 – Nov. 1**

### Major Uses:

**Pasture, Hay, Roadside Beautification**

### General Fertility Rec (pasture):

**0-50-50**

### Seasonal Production:

**November to December, and March to April**

### Tolerance:

**Soil Acidity**

### Special Notes:

**Overseeded 3-5 weeks later, avoid lime soils**

**\*Recommendations are based on regional variety test(s) from surrounding states and can be locale specific, further studying before choosing variety is best**