

Basics of Fall Vegetable Gardening

Many vegetables are well adapted to planting in the summer for fall harvest, which will extend the gardening season so you can continue to harvest fresh produce after earlier crops have finished producing. The fall harvest can be extended even further if you protect the plants from early frosts or plant them in cold frames or hotbeds.

Many cool-season vegetables, such as carrots, broccoli, cauliflower, and Brussels sprouts, produce their best flavor and quality when they mature during cool weather. In Alabama, the spring temperatures often heat up quickly making vegetables such as lettuce and spinach bolt or develop a bitter flavor when they mature during hot summer weather.

Growing a productive fall vegetable garden requires thoughtful planning and good cultural practices. In Alabama, August and September are the main planting times for a fall garden. Depending on your specific location, you may need to adjust the planting dates. For a more accurate planting schedule, determine the average date of the first killing frost in the fall, and then count backward from the frost date, using the number of days to maturity to determine the best time to plant in your area.

Preparing the Site

Before preparing the soil for a fall garden, you must decide what to do with the remains of the spring garden. In most cases, the decision is not difficult because the warm-season vegetables are beginning to look ragged. Remove the previous crop residue and any weed growth, and then till or spade the soil to a depth of at least 6 to 8 inches.

If the spring crops were heavily fertilized, you may not need to make an initial preplant fertilization. If not, you can apply 1 to 2 pounds of a complete fertilizer such as 10-10-10 per 100 square feet of bed space. Be sure to thoroughly incorporate the fertilizer.



Planting the Fall Garden

Direct seeding, which involves planting seeds rather than using transplants, for crops such as broccoli, cabbage, and collards is often used in the fall. However, the success of this planting method depends on having adequate moisture available to keep the young seedlings actively growing after germination. An overhead sprinkler can help provide seeds with sufficient moisture to germinate.

Alabama summers can be hot and dry, and soils may form a hard crust over the seeds. This crust can interfere with germination, particularly in heavy clay soil. Lettuce and spinach seeds will not germinate if the soil temperature exceeds 85 degrees F. Be sure to keep the soil moist until the young seedlings have emerged.



Watering/Fertilizing

Most vegetables require 1 inch of water per week. It is best to make a single watering that penetrates deeply rather than make frequent shallow applications. However, young seedlings and germinating seeds may need more frequent, light waterings. Do not allow seedlings to dry out. New transplants will also benefit from frequent light waterings until they develop new roots.

Many fall-maturing vegetables benefit from sidedressing with nitrogen just as spring-maturing vegetables benefit. Most leafy vegetables benefit from an application of nitrogen 3 and 6 weeks after planting.

Insects and Diseases

It is not uncommon for insects and diseases to be more abundant in the fall, mostly as a result of a buildup in their populations during the spring and summer. You may be able to keep these pests at tolerable levels, if you follow a few strategies. Strive to keep fall vegetables healthy and actively growing because healthy plants are less susceptible to insects and diseases. Check the plants frequently for insect and disease damage. If significant damage is detected, use an approved pesticide. Certain vegetables, such as squash, corn, and cucumbers, are especially insect- and disease-prone during late summer and fall.

Frost Protection

You can extend the season of tender vegetables by protecting them through the first early frost. In Alabama, we often enjoy several weeks of good growing conditions after the first frost. Cover growing beds or rows with burlap or a floating row cover supported by stakes or wire to keep the material from directly touching the plants. Protect individual plants by covering them with milk jugs, paper caps, or water-holding walls.


Most semihardy and hardy vegetables require little or no frost protection, but semihardy vegetables should be harvested before a heavy freeze, and root crops such as carrots and radishes should be harvested or mulched heavily before a hard freeze. Mulched root crops can often be harvested well into the winter, and during mild winters, harvest may continue until spring.

Suggested Fall Vegetable Planting Guide

Vegetables	Suggested Planting ¹	Specific Characteristics	Inches Between Plants	Planting Depth (inches)	Cold Tolerance	Days to Maturity
Beets	July 15 to Aug. 15	Ruby Queen, Early Wonder, Red Ace, Pacemaker II	2	0.5 to 1.0	Semi-hardy	55 to 60
Broccoli	July 15 to Aug. 15	DeCiccio, Packman, Premium Crop, Green Duke, Emperor	18	0.5 to 1.0	Hardy	70 to 80
Brussels sprouts	July 1 to 15	Long Island Improved, Jade Cross Hybrid	20	0.5 to 1.0	Hardy	90 to 100
Cabbage (plants)	Aug 1 to 15	Round Dutch, Early Jersey Wakefield, Red Express, Red Rookie	12	0.5 to 1.0	Hardy	70 to 80
Cabbage, Chinese	Aug. 1 to 15	Pak Choi, Mei Ching, Jade Pagoda, China Pride	12	0.5 to 1.0	Hardy	75-85
Carrots	July 1 to 15	Danvers Half Long, Spartan Bonus, Little Finger, Thumbelina, Scarlet Nantes	2	0.25 to 0.5	Hardy	85 to 95
Cauliflower	Aug 1 to 15	Early Snowball "A", Violet Queen, Snowcrown	18	0.5 to 1.0	Semi-hardy	55 to 65
Collards	July 15 to Aug. 15	Vates, Morris' Improved Heading, Carolina, Blue Max	18	0.5 to 1.0	Hardy	60 to 100
Cucumbers	Aug. 1 to 15	Poinsett 76, Sweet Slice, County Fair '83, Salad Bush, Fanfare	10	1.0 to 1.5	Tender	40 to 50
Kale	Aug. 15 to Sept. 1	Green Curled Scotch, Early Siberian, Vates, Dwarf Blue Curled Scotch, Blue Knight	6	0.5 to 1.0	Hardy	40 to 50
Kohlrabi	Aug. 1 to Sept. 1	White Vienna, Grand Duke Hybrid	4	0.5 to 1.0	Hardy	50 to 60
Lettuce (leaf)	Aug. 1 to Sept. 1	Grand Rapids, Salad Bowl, Buttercrunch, Red Sails, Romulus	6	0.25 to 0.5	Semi-hardy	40 to 50
Onions (seeds)	Sept. 1 to 30	Texas 1015, Granex 33, Candy	4	0.5 to 1.0	Hardy	130 to 150
Onions (sets or plants)	Sept. 1 to 15	Ebenezer, Early Grano	4	--	Hardy	60 to 80
Radishes	Aug. 15 to Sept. 15	Early Scarlet Globe, Cherry Belle, Snowbells, White Icicle	1	0.5 to 1.0	Hardy	25 to 30
Radish, Diakon	Aug. 15 to Sept. 15	April Cross, H. N. Cross	4	0.5 to 1.0	Hardy	60 to 75
Rutabagas	July 1 to Aug. 1	American Purple Top, Laurentian	4	0.5 to 1.0	Semi-hardy	70 to 80
Spinach	Aug. 1 to 15	Hybrid #7, Dark Green Bloomsdale, Tye Hybrid	6	0.5 to 1.0	Hardy	50 to 60
Turnips	Aug. 1 to 31	Purple Top White Globe, Just Right, Tokyo Cross Hybrid, White Egg, All Top	2	0.5 to 1.0	Hardy	55 to 60

¹ In north Alabama, planting dates should be approximately 2 weeks earlier in the fall. For south Alabama, planting dates should be approximately 2 weeks later in the fall.

ARCHIVE



ANR-1422

J.M. Kemble, *Extension Vegetable Specialist*, Professor, Horticulture, Auburn University.

For more information, call your county Extension office. Look in your telephone directory under your county's name to find the number.

Published by the Alabama Cooperative Extension System (Alabama A&M University and Auburn University), an equal opportunity educator and employer.

Revised Aug 2013, ANR-1422

© 2013 by the Alabama Cooperative Extension System. All rights reserved.
www.aces.edu